

9 Parametric Equations and Polar Coordinates

Dr. Jeongho Ahn

Department of Mathematics & Statistics

ASU

Outline of Chapter 9

- ① **Parametric Equations**
- ② **Calculus and Parametric equations**
- ③ **Polar Coordinates**
- ④ Calculus in Polar Coordinates
- ⑤ Conic Sections in Polar Coordinates

9.1 Parametric Equations

- When you consider a particle which moves along a curve, considering **parametric equations** is a better way to describe the curve.
- We suppose that x and y are both given as functions of a third variable t (parameter) by the **Parametric Equations**

$$x = f(t), \quad y = g(t)$$

Each parameter t determines a point $(x, y) = (f(t), g(t))$ which traces out a parameter curve C as t varies.

Example1

The following parametric equations

$$x(t) = t^2 - 3t \quad y(t) = t - 1 \quad \text{for all real numbers } t$$

will be the parabola (Cartesian equation) $x = y^2 - y - 2$.

- We can restrict t ($a \leq t \leq b$) to consider parametric equations

$$x = f(t) \quad y = g(t) \quad a \leq t \leq b,$$

where $(f(a), g(a))$ is an initial point and $(f(b), g(b))$ is a terminal point.

Example2

1. Find the curve represented by the following parametric equations

$$x = \cos t \quad y = \sin t \quad 0 \leq t \leq 2\pi.$$

2. Find the curve represented by the following parametric equations

$$x = 3 \cos 4t \quad y = 3 \sin 4t \quad 0 \leq t \leq 2\pi.$$

3. We can see that $(x - h)^2 + (y - k)^2 = r^2$ (the equation of circles) is equivalent to the following parametric equations

$$x = h + r \cos t \quad y = k + r \sin t \quad 0 \leq t \leq 2\pi.$$

Example3

Sketch the curve with the parametric equations

$$x = \cos t \quad y = \cos^2 t$$

Example4

Eliminate the parameter to find a Cartesian equation of the curve.

1. $x = 5t - 3 \quad y = 2t - 1$

2. $x = \sqrt{t} \quad y = t - 1$

3. $x = e^{3t} \quad y = t + 2 \quad t \geq -2$

Example5

Describe the motion of a particle with position (x, y) as t varies in the given interval

$$x = 1 + 2\cos t \quad y = -2 + 2\sin t, \quad \pi/2 \leq t \leq 3\pi/2$$