

## 11.4 The Cross Product

- The cross product is defined in the 3-dimensional system and is used to describe how a plane is tilting.
- The cross product has too many applications of engineering and physics.

### Fact

1. How to find the **determinant of  $3 \times 3$  matrix**?

$$\begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} = a_{11} \begin{vmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{vmatrix} - a_{12} \begin{vmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{vmatrix} + a_{13} \begin{vmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{vmatrix}.$$

2. How to find the **determinant of  $2 \times 2$  matrix**?

$$\begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - bc.$$

- For  $u = \langle u_1, u_2, u_3 \rangle$  and  $v = \langle v_1, v_2, v_3 \rangle$ , we can express  $u \times v$  in the expansion of the symbolic determinant:

$$u \times v = \begin{vmatrix} i & j & k \\ u_1 & u_2 & u_3 \\ v_1 & v_2 & v_3 \end{vmatrix} = \begin{vmatrix} u_2 & u_3 \\ v_2 & v_3 \end{vmatrix} i - \begin{vmatrix} u_1 & u_3 \\ v_1 & v_3 \end{vmatrix} j + \begin{vmatrix} u_1 & u_2 \\ v_1 & v_2 \end{vmatrix} k.$$

### Example 1

Find  $u \times v$  and  $v \times u$  if  $u = 2i - j + k$  and  $v = -4i + 3j - k$ .

### Definition

$$u \times v = (|u||v|\sin \theta)n.$$

- If  $u \neq 0$  and  $v \neq 0$  are not parallel, they determine a plane. Then we select a unit vector  $n$  perpendicular to the plane by the right-hand rule, which implies that the vector  $u \times v$  is perpendicular to both  $u$  and  $v$ .

## Theorem

*The vector  $u \times v \perp u$  and  $v$ .*

- The Theorem can be proved by using  $a \cdot b = 0 \Leftrightarrow a \perp b$ .

## Corollary

*Parallel Vectors*

$$u // v \Leftrightarrow u \times v = 0.$$

## Fact

*$|u \times v|$  is the area of a parallelogram.*

- It follows from the definition of the cross product that

$$|u \times v| = |u||v||\sin \theta|n = |u||v|\sin \theta$$

## Example2

1. Find a vector perpendicular to the plane of  $P(-1,1,0)$ ,  $Q(2,-1,1)$ , and  $R(-1,1,2)$ . Note that the vector is not unique.
2. Find the area of the triangle with vertices  $P$ ,  $Q$ , and  $R$ .

- For the standard unit vectors  $i, j, k$  we can see  
 $i \times j = k, \quad j \times k = i, \quad k \times i = j$  and  
 $i \times i = 0, \quad j \times j = 0, \quad k \times k = 0.$

- **Properties of the Cross Product**

- 1  $v \times u = -(u \times v)$
  - 2  $(ru) \times v = r(u \times v) = u \times (rv)$
  - 3  $u \times (v + w) = u \times v + u \times w$
  - 4  $(v + w) \times u = v \times u + w \times u$
  - 5  $u \cdot (v \times w) = (u \times v) \cdot w$
  - 6  $u \times (v \times w) = (u \cdot w)u - (u \cdot v)w$
  - 7  $0 \times u = 0$
- Those properties can be shown easily by writing the component form of the vectors and using the symbolic expression.
  - Note that  $(u \times v) \times w \neq u \times (v \times w)$

## Definition

Triple Scalar or Box Product  $u \cdot (v \times w)$

$|u \cdot (v \times w)|$  is the volume of the parallelepiped, since

$$|u \cdot (v \times w)| = |u||v \times w| |\cos \theta|.$$

- The triple scalar product can be calculated as the determinant:

$$u \cdot (v \times w) = \begin{vmatrix} u_1 & u_2 & u_3 \\ v_1 & v_2 & v_3 \\ w_1 & w_2 & w_3 \end{vmatrix}.$$

- By using the determinant, we can see that  
 $(u \times v) \cdot w = (v \times w) \cdot u = (w \times u) \cdot v.$

## Example3

Find the volume of the box(parallelepiped) determined by  
 $u = i - 2j - k$ ,  $v = -2i + k$  and  $w = 5j - 4k$ .